arXiv:1006.1561v1 [math.CA] 8 Jun 2010

NEW OSTROWSKI TYPE INEQUALITIES FOR $m-$CONVEX FUNCTIONS AND APPLICATIONS

HAVVA KAVURMACI[▲,★], M. EMİN ÖZDEMİR[♠], AND MERVE AVCI[▲]

Abstract. In this paper, we establish new inequalities of Ostrowski type for functions whose derivatives in absolute value are $m-$convex. We also give some applications to special means of positive real numbers. Finally, we obtain some error estimates for the midpoint formula.

1. INTRODUCTION

Let $f : I \subset [0,\infty) \to \mathbb{R}$ be a differentiable mapping on I°, the interior of the interval I, such that $f' \in L([a,b])$ where a , $b \in I$ with $a < b$. If $|f'(x)| \leq M$, then the following inequality holds (see [2]):

$$\left| f(x) - \frac{1}{b-a}\int_a^b f(u)du \right| \leq \frac{M}{b-a}\left[\frac{(x-a)^2+(b-x)^2}{2}\right].$$

This inequality is well known in the literature as the *Ostrowski inequality.* For some results which generalize, improve, and extend the above inequality, see [2],[5],[6],[10] and [11], the references therein.

In [12], G. Toader defined $m-$convexity, an intermediate between the usual convexity and starshaped property, as the following:

Definition 1. *The function $f : [0,b] \to \mathbb{R}$, $b > 0$, is said to be $m-$convex, where $m \in [0,1]$, if we have*

$$f(tx + m(1-t)y) \leq tf(x) + m(1-t)f(y)$$

for all $x, y \in [0,b]$ and $t \in [0,1]$.

Denote by $K_m(b)$ the set of the $m-$convex functions on $[0,b]$ for which $f(0) \leq 0$.

Definition 2. *The function $f : [0,b] \to \mathbb{R}$, $b > 0$ is said to be starshaped if for every $x \in [0,b]$ and $t \in [0,1]$ we have:*

$$f(tx) \leq tf(x).$$

For $m = 1$, we recapture the concept of convex functions defined on $[0,b]$ and $m = 0$ we get the concept of starshaped functions on $[0,b]$.

The following theorem contains the Hermite-Hadamard type integral inequality (see [8]).

Theorem 1. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be an M-Lipschitzian mapping on I and $a, b \in I$ with $a < b$. Then we have the inequality:*

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a}\int_a^b f(x)dx \right| \leq M\frac{(b-a)}{4}. \tag{1}$$

In [13] E. Set, M.E. Özdemir, M.Z. Sarıkaya established the following theorem.

Theorem 2. *Let $f : I^{\circ} \subset [0,b^*] \to \mathbb{R}$, $b^* > 0$, be a differentiable mapping on I°, $a, b \in I^{\circ}$ with $a < b$. If $|f'|^q$ is $m-$convex on $[a,b]$, $q > 1$ and $m \in (0,1]$, then the following inequality holds:*

2000 *Mathematics Subject Classification.* 26A51, 26D10, 26D15.

Key words and phrases. $m-$convex function, Starshaped function, Convex function, Ostrowski inequality, Hermite-Hadamard inequality, Hölder inequality, Power Mean inequality, Special means, The Midpoint formula, Lipschitzian mapping.

[★] Corresponding Author.

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right|\leq (b-a)\left(\frac{3^{1-\left(\frac{1}{q}\right)}}{8}\right)\left(\left|f'(a)\right|+m^{\frac{1}{q}}\left|f'(\frac{b}{m})\right|\right). \tag{2}$$

where $\frac{b}{m}<b^*$.

In [14] U. Kirmaci proved the following theorem.

Theorem 3. *Let $f:I^\circ\subset\mathbb{R}\to\mathbb{R}$ be a differentiable mapping on I°, $a,b\in I^\circ$ with $a<b$. If the mapping $|f'|$ is convex on $[a,b]$, then we have*

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right|\leq \frac{b-a}{8}\left(\left|f'(a)\right|+\left|f'(b)\right|\right). \tag{3}$$

In [9] S.S. Dragomir and G. Toader proved the following Hermite-Hadamard type inequality for $m-$convex functions.

Theorem 4. *Let $f:[0,\infty)\to\mathbb{R}$ be an $m-$convex function with $m\in(0,1]$. If $0\leq a<b<\infty$ and $f\in L^1([a,b])$ then*

$$\frac{1}{b-a}\int_a^b f(x)dx\leq \min\left\{\frac{f(a)+mf(\frac{b}{m})}{2},\frac{f(b)+mf(\frac{a}{m})}{2}\right\}. \tag{4}$$

Some generalizations of this result can be found in [4].

In [3] M.K. Bakula, M.E. Özdemir and J. Pečarić proved the following theorems.

Theorem 5. *Let I be an open real interval such that $[0,\infty)\subset I$. Let $f:I\to\mathbb{R}$ be a differentiable function on I such that $f'\in L([a,b])$, where $0\leq a<b<\infty$. If $|f'|^q$ is $m-$convex on $[a,b]$ for some fixed $m\in(0,1]$ and $q\in[1,\infty)$, then*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right|\leq \frac{b-a}{4}\left(\mu_1^{\frac{1}{q}}+\mu_2^{\frac{1}{q}}\right), \tag{5}$$

where

$$\begin{aligned}\mu_1 &= \min\left\{\frac{|f'(a)|^q+m\left|f'(\frac{a+b}{2m})\right|^q}{2},\frac{\left|f'(\frac{a+b}{2})\right|^q+m\left|f'(\frac{a}{m})\right|^q}{2}\right\},\\ \mu_2 &= \min\left\{\frac{|f'(b)|^q+m\left|f'(\frac{a+b}{2m})\right|^q}{2},\frac{\left|f'(\frac{a+b}{2})\right|^q+m\left|f'(\frac{b}{m})\right|^q}{2}\right\}.\end{aligned}$$

Theorem 6. *Let I be an open real interval such that $[0,\infty)\subset I$. Let $f:I\to\mathbb{R}$ be a differentiable function on I such that $f'\in L([a,b])$, where $0\leq a<b<\infty$. If $|f'|^q$ is $m-$convex on $[a,b]$ for some fixed $m\in(0,1]$ and $q\in[1,\infty)$, then*

$$\begin{aligned}&\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right| \\ \leq\ & \frac{b-a}{4}\min\left\{\left(\frac{|f'(a)|^q+m\left|f'(\frac{b}{m})\right|^q}{2}\right)^{\frac{1}{q}},\left(\frac{m\left|f'(\frac{a}{m})\right|^q+|f'(b)|^q}{2}\right)^{\frac{1}{q}}\right\}.\end{aligned} \tag{6}$$

The main purpose of this paper is to establish new Ostrowski type inequalities for functions whose derivatives in absolute value are $m-$convex. Using these results we give some applications to special means of positive real numbers and we obtain some error estimates for the midpoint formula.

2. THE RESULTS

In [1], in order to prove some inequalities related to Ostrowski inequality, M. Alomari and M. Darus used the following lemma with the constant $(b-a)$, but we changed it with the constant $(a-b)$ to obtain an equality in Lemma 7.

Lemma 7. *Let $f:I\subset\mathbb{R}\rightarrow\mathbb{R}$ be a differentiable mapping on I° where $a,b\in I$ with $a<b$. If $f'\in L\left([a,b]\right)$, then the following equality holds:*

$$f(x)-\frac{1}{b-a}\int_a^b f(u)du=(a-b)\int_0^1 p(t)f'(ta+(1-t)b)dt \tag{7}$$

for each $t\in[0,1]$, where

$$p(t)=\begin{cases} t & , \quad t\in\left[0,\frac{b-x}{b-a}\right] \\ t-1 & , \quad t\in\left(\frac{b-x}{b-a},1\right] \end{cases},$$

for all $x\in[a,b]$.

Theorem 8. *Let I be an open real interval such that $[0,\infty)\subset I$. Let $f:I\rightarrow\mathbb{R}$ be a differentiable function on I such that $f'\in L([a,b])$, where $0\leq a<b<\infty$. If $|f'|$ is $m-$convex on $[a,b]$ for some fixed $m\in(0,1]$, then the following inequality holds:*

$$\begin{aligned} &\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right| \\ &\leq (b-a)\min\left\{\begin{array}{c} \left[\frac{1}{6}-\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2+\frac{2}{3}\left(\frac{b-x}{b-a}\right)^3\right]|f'(a)| \\ +m\left[\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2-\frac{1}{3}\left(\frac{b-x}{b-a}\right)^3+\frac{1}{3}\left(\frac{x-a}{b-a}\right)^3\right]\left|f'\left(\frac{b}{m}\right)\right| \end{array}\right., \\ &\qquad\left.\begin{array}{c} \left[\frac{1}{6}-\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2+\frac{2}{3}\left(\frac{b-x}{b-a}\right)^3\right]|f'(b)| \\ +m\left[\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2-\frac{1}{3}\left(\frac{b-x}{b-a}\right)^3+\frac{1}{3}\left(\frac{x-a}{b-a}\right)^3\right]\left|f'\left(\frac{a}{m}\right)\right| \end{array}\right\}. \end{aligned} \tag{8}$$

for each $x\in[a,b]$.

Proof. By Lemma 7, we have

$$\begin{aligned} &\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right| \\ \leq\ & (b-a)\int_0^{\frac{b-x}{b-a}} t\left|f'(ta+(1-t)b)\right|dt \\ &+(b-a)\int_{\frac{b-x}{b-a}}^1 (1-t)\left|f'(ta+(1-t)b)\right|dt \end{aligned}$$

Since $|f'|$ is $m-$convex on $[a,b]$ we know that for any $t\in[0,1]$

$$\begin{aligned} \left|f'(ta+(1-t)b)\right| &= \left|f'(ta+m(1-t)\frac{b}{m})\right| \\ &\leq t\left|f'(a)\right|+m(1-t)\left|f'\left(\frac{b}{m}\right)\right|, \end{aligned}$$

Hence

$$\begin{aligned}
&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right| \\
\leq\ & (b-a)\int_0^{\frac{b-x}{b-a}} t\left[t\left|f'(a)\right|+m(1-t)\left|f'\left(\frac{b}{m}\right)\right|\right]dt \\
&+(b-a)\int_{\frac{b-x}{b-a}}^1 (1-t)\left[t\left|f'(a)\right|+m(1-t)\left|f'\left(\frac{b}{m}\right)\right|\right]dt \\
=\ & (b-a)\left\{\begin{array}{c} \left[\frac{1}{6}-\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2+\frac{2}{3}\left(\frac{b-x}{b-a}\right)^3\right]\left|f'(a)\right| \\ +m\left[\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2-\frac{1}{3}\left(\frac{b-x}{b-a}\right)^3+\frac{1}{3}\left(\frac{x-a}{b-a}\right)^3\right]\left|f'\left(\frac{b}{m}\right)\right| \end{array}\right\}
\end{aligned}$$

where we use the facts that

$$\begin{aligned}
&\int_0^{\frac{b-x}{b-a}} t\left[t\left|f'(a)\right|+m(1-t)\left|f'\left(\frac{b}{m}\right)\right|\right]dt \\
=\ & \frac{1}{3}\left(\frac{b-x}{b-a}\right)^3\left|f'(a)\right|+m\left[\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2-\frac{1}{3}\left(\frac{b-x}{b-a}\right)^3\right]\left|f'\left(\frac{b}{m}\right)\right|,
\end{aligned}$$

and

$$\begin{aligned}
&\int_{\frac{b-x}{b-a}}^1 (1-t)\left[t\left|f'(a)\right|+m(1-t)\left|f'\left(\frac{b}{m}\right)\right|\right]dt \\
=\ & \left[\frac{1}{6}-\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2+\frac{1}{3}\left(\frac{b-x}{b-a}\right)^3\right]\left|f'(a)\right|+m\frac{1}{3}\left(\frac{x-a}{b-a}\right)^3\left|f'\left(\frac{b}{m}\right)\right|,
\end{aligned}$$

and analogously

$$\begin{aligned}
&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right| \\
\leq\ & (b-a)\left\{\begin{array}{c} \left[\frac{1}{6}-\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2+\frac{2}{3}\left(\frac{b-x}{b-a}\right)^3\right]\left|f'(b)\right| \\ +m\left[\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2-\frac{1}{3}\left(\frac{b-x}{b-a}\right)^3+\frac{1}{3}\left(\frac{x-a}{b-a}\right)^3\right]\left|f'\left(\frac{a}{m}\right)\right| \end{array}\right\}.
\end{aligned}$$

The proof is completed. □

Remark 1. *Suppose that all the assumptions of Theorem 8 are satisfied. If we choose* $x=\frac{a+b}{2}$, *then we have*

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(u)du\right| \\
\leq\ & \frac{b-a}{8}\min\left\{\left|f'(a)\right|+m\left|f'\left(\frac{b}{m}\right)\right|,\left|f'(b)\right|+m\left|f'\left(\frac{a}{m}\right)\right|\right\}
\end{aligned}$$

which is [(6),q=1].

Remark 2. *Suppose that all the assumptions of Theorem 8 are satisfied. Then*

(A) If we choose $m=1$ and $x=\frac{a+b}{2}$, we obtain

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(u)du\right|\leq\frac{b-a}{8}\left(\left|f'(a)\right|+\left|f'(b)\right|\right),$$

which is (3).

(B) In (A). Additionally, if we choose $|f'(x)|\leq M$, $M>0$

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(u)du\right|\leq M\frac{(b-a)}{4}$$

which is (1).

Theorem 9. *Let I be an open real interval such that $[0,\infty)\subset I$. Let $f:I\to\mathbb{R}$ be a differentiable function on I such that $f'\in L([a,b])$, where $0\leq a<b<\infty$. If $|f'|^{\frac{p}{p-1}}$ is $m-$convex on $[a,b]$ for some fixed $m\in(0,1]$ and $p>1$, $\frac{1}{p}+\frac{1}{q}=1$, then the following inequality holds:*

$$\tag{9}\begin{aligned}&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right|\\ &\leq\frac{1}{(p+1)^{\frac{1}{p}}}\left\{\begin{array}{c}\frac{(b-x)^2}{b-a}\left[\min\left\{\frac{|f'(b)|^q+m\left|f'\left(\frac{x}{m}\right)\right|^q}{2},\frac{|f'(x)|^q+m\left|f'\left(\frac{b}{m}\right)\right|^q}{2}\right\}\right]^{\frac{1}{q}}\\ +\frac{(x-a)^2}{b-a}\left[\min\left\{\frac{|f'(a)|^q+m\left|f'\left(\frac{x}{m}\right)\right|^q}{2},\frac{|f'(x)|^q+m\left|f'\left(\frac{a}{m}\right)\right|^q}{2}\right\}\right]^{\frac{1}{q}}\end{array}\right\}\end{aligned}$$

for each $x\in[a,b]$.

Proof. From Lemma 7 and using the Hölder inequality, we have

$$\begin{aligned}&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right|\\ \leq\ &(b-a)\left(\int_0^{\frac{b-x}{b-a}}t^p dt\right)^{\frac{1}{p}}\left(\int_0^{\frac{b-x}{b-a}}\left|f'\left(ta+(1-t)b\right)\right|^q dt\right)^{\frac{1}{q}}\\ &+(b-a)\left(\int_{\frac{b-x}{b-a}}^1(1-t)^p dt\right)^{\frac{1}{p}}\left(\int_{\frac{b-x}{b-a}}^1\left|f'\left(ta+(1-t)b\right)\right|^q dt\right)^{\frac{1}{q}}\\ \leq\ &(b-a)\left(\frac{b-x}{b-a}\right)^{\frac{p+1}{p}}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{b-x}{b-a}\right)^{\frac{1}{q}}\\ &\times\left(\min\left\{\frac{|f'(b)|^q+m\left|f'\left(\frac{x}{m}\right)\right|^q}{2},\frac{|f'(x)|^q+m\left|f'\left(\frac{b}{m}\right)\right|^q}{2}\right\}\right)^{\frac{1}{q}}\\ &+(b-a)\left(\frac{x-a}{b-a}\right)^{\frac{p+1}{p}}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{x-a}{b-a}\right)^{\frac{1}{q}}\\ &\times\left(\min\left\{\frac{|f'(a)|^q+m\left|f'\left(\frac{x}{m}\right)\right|^q}{2},\frac{|f'(x)|^q+m\left|f'\left(\frac{a}{m}\right)\right|^q}{2}\right\}\right)^{\frac{1}{q}}\\ =\ &\frac{1}{(p+1)^{\frac{1}{p}}}\frac{1}{b-a}\left\{\begin{array}{c}(b-x)^2\left[\min\left\{\frac{|f'(b)|^q+m\left|f'\left(\frac{x}{m}\right)\right|^q}{2},\frac{|f'(x)|^q+m\left|f'\left(\frac{b}{m}\right)\right|^q}{2}\right\}\right]^{\frac{1}{q}}\\ +(x-a)^2\left[\min\left\{\frac{|f'(a)|^q+m\left|f'\left(\frac{x}{m}\right)\right|^q}{2},\frac{|f'(x)|^q+m\left|f'\left(\frac{a}{m}\right)\right|^q}{2}\right\}\right]^{\frac{1}{q}}\end{array}\right\}\end{aligned}$$

where we use the facts that

$$\begin{aligned}\int_0^{\frac{b-x}{b-a}}t^p dt&=\left(\frac{b-x}{b-a}\right)^{p+1}\frac{1}{p+1},\\ \int_{\frac{b-x}{b-a}}^1(1-t)^p dt&=\left(\frac{x-a}{b-a}\right)^{p+1}\frac{1}{p+1},\end{aligned}$$

and by Theorem 4 we get

$$\begin{aligned}
&\frac{b-a}{b-x}\int_0^{\frac{b-x}{b-a}}\left|f'(ta+(1-t)b)\right|^q dt \\
\leq\ &\min\left\{\frac{|f'(b)|^q+m\left|f'\left(\frac{x}{m}\right)\right|^q}{2},\frac{|f'(x)|^q+m\left|f'\left(\frac{b}{m}\right)\right|^q}{2}\right\}, \\
&\frac{b-a}{x-a}\int_{\frac{b-x}{b-a}}^{1}\left|f'(ta+(1-t)b)\right|^q dt \\
\leq\ &\min\left\{\frac{|f'(a)|^q+m\left|f'\left(\frac{x}{m}\right)\right|^q}{2},\frac{|f'(x)|^q+m\left|f'\left(\frac{a}{m}\right)\right|^q}{2}\right\}.
\end{aligned}$$

The proof is completed. □

Corollary 10. *Suppose that all the assumptions of Theorem 9 are satisfied, if we choose* $|f'(x)|\leq M$*,* $M>0$*, then we have*

$$\begin{aligned}
&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right| \\
\leq\ &\left(\frac{1}{(p+1)^{\frac{1}{p}}}\right)\left(\frac{1+m}{2}\right)^{\frac{1}{q}}M\left[\frac{(b-x)^2+(x-a)^2}{b-a}\right].
\end{aligned}$$

Corollary 11. *Suppose that all the assumptions of Theorem 9 are satisfied, if we choose* $x=\frac{a+b}{2}$ *and* $\frac{1}{2}<\left(\frac{1}{p+1}\right)^{\frac{1}{p}}<1$*, then we have*

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(u)du\right|\leq\frac{b-a}{4}\left(\mu_1^{\frac{1}{q}}+\mu_2^{\frac{1}{q}}\right),$$

where

$$\begin{aligned}
\mu_1 &= \min\left\{\frac{|f'(b)|^q+m\left|f'(\frac{a+b}{2m})\right|^q}{2},\frac{\left|f'(\frac{a+b}{2})\right|^q+m\left|f'\left(\frac{b}{m}\right)\right|^q}{2}\right\}, \\
\mu_2 &= \min\left\{\frac{|f'(a)|^q+m\left|f'(\frac{a+b}{2m})\right|^q}{2},\frac{\left|f'(\frac{a+b}{2})\right|^q+m\left|f'(\frac{a}{m})\right|^q}{2}\right\}.
\end{aligned}$$

Remark 3. *Corollary 11 is similar to (5) inequality, but for the left-hand side of Hermite-Hadamard inequality.*

Remark 4. *Suppose that all the assumptions of Theorem 9 are satisfied. Then in Corollary 11*

(D) $|f'|$ is increasing and $m=1$ then we have

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(u)du\right|\leq\frac{b-a}{2}\left|f'(b)\right|,$$

(E) $|f'|$ is decreasing and $m=1$ then we have

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(u)du\right|\leq\frac{b-a}{2}\left|f'(a)\right|,$$

(F) $|f'(b)|=|f'(a)|=\left|f'\left(\frac{a+b}{2}\right)\right|$ and $m=1$ then we have

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(u)du\right|\leq\frac{b-a}{2}\left|f'\left(\frac{a+b}{2}\right)\right|.$$

Theorem 12. *Let I be an open real interval such that $[0,\infty)\subset I$. Let $f:I\to\mathbb{R}$ be a differentiable function on I such that $f'\in L([a,b])$, where $0\le a<b<\infty$. If $|f'|^q$ is $m-$convex on $[a,b]$ for some fixed $m\in(0,1]$ and $q\in[1,\infty)$, $x\in[a,b]$, then the following inequality holds:*

$$\begin{aligned}
&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right| \\
\le\ & (b-a)\left(\frac{1}{2}\right)^{1-\frac{1}{q}} \\
&\left\{\begin{array}{c}
\left(\frac{b-x}{b-a}\right)^{2\left(1-\frac{1}{q}\right)}\left[\frac{1}{3}\left(\frac{b-x}{b-a}\right)^{3}|f'(a)|^q+m\frac{(b-x)^2(b-3a+2x)}{6(b-a)^3}\left|f'\left(\frac{b}{m}\right)\right|^q\right]^{\frac{1}{q}} \\
+\left(\frac{x-a}{b-a}\right)^{2\left(1-\frac{1}{q}\right)}\left[\left(\frac{1}{6}+\frac{(b-x)^2(3a-b-2x)}{6(b-a)^3}\right)|f'(a)|^q+m\frac{1}{3}\left(\frac{x-a}{b-a}\right)^{3}\left|f'\left(\frac{b}{m}\right)\right|^q\right]^{\frac{1}{q}}
\end{array}\right\}
\end{aligned}\tag{10}$$

for each $x\in[a,b]$.

Proof. By Lemma 7 and using the well known power mean inequality we have

$$\begin{aligned}
&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right| \\
\le\ & (b-a)\int_0^{\frac{b-x}{b-a}} t\left|f'(ta+(1-t)b)\right|dt \\
&+(b-a)\int_{\frac{b-x}{b-a}}^1 (1-t)\left|f'(ta+(1-t)b)\right|dt \\
\le\ & (b-a)\left(\int_0^{\frac{b-x}{b-a}} tdt\right)^{1-\frac{1}{q}}\left(\int_0^{\frac{b-x}{b-a}} t\left|f'(ta+(1-t)b)\right|^q dt\right)^{\frac{1}{q}} \\
&+(b-a)\left(\int_{\frac{b-x}{b-a}}^1 (1-t)dt\right)^{1-\frac{1}{q}}\left(\int_{\frac{b-x}{b-a}}^1 (1-t)\left|f'(ta+(1-t)b)\right|^q dt\right)^{\frac{1}{q}} \\
\le\ & (b-a)\left(\frac{1}{2}\right)^{1-\frac{1}{q}} \\
&\left\{\begin{array}{c}
\left(\frac{b-x}{b-a}\right)^{2\left(1-\frac{1}{q}\right)}\left[\frac{1}{3}\left(\frac{b-x}{b-a}\right)^{3}|f'(a)|^q+m\frac{(b-x)^2(b-3a+2x)}{6(b-a)^3}\left|f'\left(\frac{b}{m}\right)\right|^q\right]^{\frac{1}{q}} \\
+\left(\frac{x-a}{b-a}\right)^{2\left(1-\frac{1}{q}\right)}\left[\left(\frac{1}{6}+\frac{(b-x)^2(3a-b-2x)}{6(b-a)^3}\right)|f'(a)|^q+m\frac{1}{3}\left(\frac{x-a}{b-a}\right)^{3}\left|f'\left(\frac{b}{m}\right)\right|^q\right]^{\frac{1}{q}}
\end{array}\right\}
\end{aligned}$$

where we use the facts that

$$\int_0^{\frac{b-x}{b-a}} tdt=\frac{1}{2}\left(\frac{b-x}{b-a}\right)^2,$$

$$\begin{aligned}
&\int_0^{\frac{b-x}{b-a}} t\left|f'(ta+(1-t)b)\right|^q dt \\
\le\ & \frac{1}{3}\left(\frac{b-x}{b-a}\right)^3|f'(a)|^q+m\frac{(b-x)^2(b-3a+2x)}{6(b-a)^3}\left|f'\left(\frac{b}{m}\right)\right|^q,
\end{aligned}$$

$$\int_{\frac{b-x}{b-a}}^1 (1-t)dt=\frac{1}{2}\left(\frac{x-a}{b-a}\right)^2,$$

$$\begin{aligned}
&\int_{\frac{b-x}{b-a}}^{1}(1-t)\left|f'(ta+(1-t)b)\right|^{q}dt\\
&\leq \left[\frac{1}{6}+\frac{(b-x)^{2}(3a-2x-b)}{6(b-a)^{3}}\right]\left|f'(a)\right|^{q}+m\frac{1}{3}\left(\frac{x-a}{b-a}\right)^{3}\left|f'\left(\frac{b}{m}\right)\right|^{q}.
\end{aligned}$$

The proof is completed. □

Remark 5. *Suppose that all the assumptions of Theorem 12 are satisfied. If we choose* $x=\frac{a+b}{2}$, *we obtain*

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_{a}^{b}f(u)du\right|\leq (b-a)\left(\frac{3^{1-\frac{1}{q}}}{8}\right)\left(\left|f'(a)\right|+m^{\frac{1}{q}}\left|f'\left(\frac{b}{m}\right)\right|\right)$$

which is (2).

3. APPLICATIONS TO SPECIAL MEANS

Let us recall the following means for two positive numbers.
(AM) *The Arithmetic mean*

$$A=A(a,b)=\frac{a+b}{2};\ a,b>0,$$

$(p-LM)$ *The p-Logarithmic mean*

$$L_{p}=L_{p}(a,b)=\begin{cases} a & if\ a=b\\ \left[\frac{b^{p+1}-a^{p+1}}{(p+1)(b-a)}\right]^{\frac{1}{p}} & if\ a\neq b\end{cases};\ a,b>0,$$

(IM) *The Identric mean*

$$I=I(a,b)=\begin{cases} a & if\ a=b\\ \frac{1}{e}\left(\frac{b^{b}}{a^{a}}\right)^{\frac{1}{b-a}} & if\ a\neq b\end{cases};\ a,b>0.$$

The following propositions hold:

Proposition 13. *Let* $a,b\in[0,\infty)$, *and* $a<b$, $n\geq 2$ *with* $m\in(0,1]$. *Then we have*

$$\left|A^{n}(a,b)-L_{n}^{n}(a,b)\right|\leq n\frac{b-a}{8}\min\left\{2A\left(a^{n-1},m\left(\frac{b}{m}\right)^{n-1}\right),2A\left((b)^{n-1},m\left(\frac{a}{m}\right)^{n-1}\right)\right\}.$$

Proof. The proof follows by Remark 1 on choosing $f:[0,\infty)\to[0,\infty)$, $f(x)=x^{n}$, $n\in\mathbb{Z},n\geq 2$ which is $m-$convex on $[0,\infty)$. □

Proposition 14. *Let* $a,b\in[0,\infty)$, *and* $a<b$, *with* $m\in(0,1]$. *Then we have*

$$\left|\ln\frac{I(a+1,b+1)}{A(a,b)+1}\right|\leq\frac{b-a}{4}\left(\eta_{1}^{\frac{1}{q}}+\eta_{2}^{\frac{1}{q}}\right),$$

where

$$\begin{aligned}
\eta_{1}^{\frac{1}{q}} &= \min\left\{\frac{\left(\frac{1}{b+1}\right)^{q}+m\left(\frac{2m}{a+b+2m}\right)^{q}}{2},\frac{\left(\frac{2}{a+b+2}\right)^{q}+m\left(\frac{m}{b+m}\right)^{q}}{2}\right\},\\
\eta_{2}^{\frac{1}{q}} &= \min\left\{\frac{\left(\frac{1}{a+1}\right)^{q}+m\left(\frac{2m}{a+b+2m}\right)^{q}}{2},\frac{\left(\frac{2}{a+b+2}\right)^{q}+m\left(\frac{m}{a+m}\right)^{q}}{2}\right\}.
\end{aligned}$$

Proof. The proof follows by Corollary 11 on choosing $f:[0,\infty)\to(-\infty,0]$, $f(x)=-\ln(x+1)$ which is $m-$convex on $[0,\infty)$, $p>1$. □

4. APPLICATIONS TO THE MIDPOINT FORMULA FOR $1-CONVEX$ FUNCTIONS

Let d be a division $a = x_0 < x_1 < ... < x_{n-1} < x_n = b$ of the interval $[a, b]$ and consider the quadrature formula

$$\int_a^b f(x)dx = M(f,d) + E(f,d), \tag{11}$$

where

$$M(f,d) = \sum_{i=1}^{n-1} (x_{i+1} - x_i) f\left(\frac{x_{i+1} + x_i}{2}\right)$$

is the midpoint formula and $E(f, d)$ denotes the associated approximation error (see [7]).

Here, we obtain some error estimates for the midpoint formula.

Proposition 15. *Let I be an open real interval such that $[0, \infty) \subset I$. Let $f : I \to \mathbb{R}$ be a differentiable function on I such that $f' \in L([a, b])$, where $0 \le a < b < \infty$. If $|f'|^q$ is $1-$convex on $[a, b]$ for some fixed $m \in (0, 1]$ and $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, then in (11), for every division d of $[a, b]$, the midpoint error satisfies*

$$|E(f,d)| \le \frac{1}{4} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 \left(\mu_1^{\frac{1}{q}} + \mu_2^{\frac{1}{q}}\right),$$

where

$$\begin{aligned}
\mu_1 &= \min\left\{\frac{|f'(x_i)|^q + \left|f'(\frac{x_i+x_{i+1}}{2})\right|^q}{2}, \frac{\left|f'(\frac{x_i+x_{i+1}}{2})\right|^q + |f'(x_i)|^q}{2}\right\} = \frac{\left|f'(\frac{x_i+x_{i+1}}{2})\right|^q + |f'(x_i)|^q}{2}, \\
\mu_2 &= \min\left\{\frac{|f'(x_{i+1})|^q + \left|f'(\frac{x_i+x_{i+1}}{2})\right|^q}{2}, \frac{\left|f'(\frac{x_i+x_{i+1}}{2})\right|^q + |f'(x_{i+1})|^q}{2}\right\} = \frac{\left|f'(\frac{x_i+x_{i+1}}{2})\right|^q + |f'(x_{i+1})|^q}{2}.
\end{aligned}$$

Proof. On applying Corollary 11 with $m = 1$ on the subinterval $[x_i, x_{i+1}]$ $(i = 0, 1, 2, ..., n-1)$ of the division, we have

$$\left|f\left(\frac{x_{i+1} + x_i}{2}\right) - \frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} f(x)dx\right| \le \frac{x_{i+1} - x_i}{4} \left(\mu_1^{\frac{1}{q}} + \mu_2^{\frac{1}{q}}\right),$$

where

$$\begin{aligned}
\mu_1 &= \frac{\left|f'(\frac{x_i+x_{i+1}}{2})\right|^q + |f'(x_i)|^q}{2} \\
\mu_2 &= \frac{\left|f'(\frac{x_i+x_{i+1}}{2})\right|^q + |f'(x_{i+1})|^q}{2}.
\end{aligned}$$

Hence, in (11) we have

$$\begin{aligned}
&\left|\int_a^b f(x)dx - M(f,d)\right| \\
&= \left|\sum_{i=0}^{n-1} \left[\int_{x_i}^{x_{i+1}} f(x)dx - (x_{i+1} - x_i) f\left(\frac{x_{i+1} - x_i}{2}\right)\right]\right| \\
&\le \sum_{i=0}^{n-1} \left|\int_{x_i}^{x_{i+1}} f(x)dx - (x_{i+1} - x_i) f\left(\frac{x_{i+1} - x_i}{2}\right)\right| \\
&\le \frac{1}{4} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 \left(\mu_1^{\frac{1}{q}} + \mu_2^{\frac{1}{q}}\right),
\end{aligned}$$

which completes the proof. ☐

Proposition 16. *Let I be an open real interval such that $[0,\infty)\subset I$. Let $f:I\rightarrow\mathbb{R}$ be a differentiable function on I such that $f'\in L([a,b])$, where $0\leq a<b<\infty$. If $|f'|^q$ is $1-$convex on $[a,b]$ for some fixed $m\in(0,1]$ and $q\in[1,\infty)$, $x\in[a,b]$, then in (11), for every division d of $[a,b]$, the midpoint error satisfies*

$$|E(f,d)|\leq\left(\frac{3^{1-\frac{1}{q}}}{8}\right)\sum_{i=0}^{n-1}(x_{i+1}-x_i)^2\left(\left|f'(x_i)\right|+\left|f'\left(x_{i+1}\right)\right|\right).$$

Proof. The proof is similar to that of Proposition 15 and using Remark 5 with $m=1$. ☐

References

[1] M. Alomari and M. Darus, Some Ostrowski's type inequalities for convex functions with applications, *RGMIA Res. Rep. Coll.,* 13(2) (2010), Article 3.

[2] M. Alomari, M. Darus, S.S. Dragomir and P. Cerone, Ostrowski's inequalities for functions whose derivatives are $s-$convex in the second sense, *RGMIA Res. Rep. Coll.,* 12(2009), Supplement, Article 15.

[3] M.K. Bakula, M.E. Özdemir and J. Pečarić, Hadamard type inequalities for $m-$convex and $(\alpha,m)-$convex functions, *J. Inequal. Pure & Appl. Math.,* 9(2008), Article 96.

[4] M.K. Bakula, J. Pečarić, and M. Ribičić, Companion inequalities to Jensen's inequality for $m-$convex and $(\alpha,m)-$convex functions, *J. Inequal. Pure & Appl. Math.,* 7(2006), Article 194.

[5] N.S. Barnett, P. Cerone, S.S. Dragomir, M.R. Pinheiro and A. Sofo, Ostrowski type inequalities for functions whose modulus of derivatives are convex and applications, *RGMIA Res. Rep. Coll.,* 5(2) (2002), Article 1.

[6] P. Cerone, S.S. Dragomir and J. Roumeliotis, An inequality of Ostrowski type for mappings whose second derivatives are bounded and applications, *RGMIA Res. Rep. Coll.,* 1(1) (1998), Article 4.

[7] C.E.M. Pearce and J. Pečarić, Inequalities for differentiable mappings with application to special means and quadrature formula, *Appl. Math. Lett.* 13(2000) 51-55.

[8] S.S. Dragomir, Y.J. Cho and S.S. Kim, Inequalities of Hadamard's type for Lipschitzian mappings and their applications, *J. of Math. Anal. Appl.,* 245(2) (2000), 489-501.

[9] S.S. Dragomir and G. Toader, Some inequalities for $m-$convex functions, *Studia Univ. Babeş-Bolyai Math.,* 38(1) (1993), 21-28.

[10] S.S. Dragomir and A. Sofo, Ostrowski type inequalities for functions whose derivatives are convex, Proceedings of the 4th International Conference on Modelling and Simulation. November 11-13, 2002. Victoria University, Melbourne, Australia. *RGMIA Res. Rep. Coll.,* 5(2002), Supplement, Article 30.

[11] S.S. Dragomir and S.Wang, Applications of Ostrowski's inequality to the estimation of error bounds for some special means and some numerical quadrature rules, *Appl. Math. Lett.,* 11(1998), 105-109.

[12] G. Toader, Some generalizations of the convexity, *Proceedings of The Colloquium On Approximation and Optimization,* Univ. Cluj-Napoca, Cluj-Napoca, 1984, 329-338.

[13] E. Set, M.E. Özdemir and M.Z. Sarıkaya, Inequalities of Hermite-Hadamard's type for functions whose derivatives absolute values are $m-$convex, *RGMIA Res. Rep. Coll.,* 13(2010), Supplement, Article 5.

[14] U.S. Kirmaci, Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, *Appl. Math. Comput., 147(2004), 137-146.*

♠ATATURK UNIVERSITY, K.K. EDUCATION FACULTY, DEPARTMENT OF MATHEMATICS, 25240, CAMPUS, ERZURUM, TURKEY
E-mail address: `havva.kvrmc@yahoo.com`

♣GRADUATE SCHOOL OF NATURAL AND APPLIED SCIENCE, AĞRI İBRAHİM ÇEÇEN UNIVERSITY, AĞRI, TURKEY
E-mail address: `emos@atauni.edu.tr`

E-mail address: `merveavci@ymail.com`